


38. (Amended) The method of claim 32 where the cancerostatic or immunosuppressive agent is selected from the group consisting of compounds of formula I:



where:

each of $R^{1(i)}$, $R^{2(i)}$, $R^{3(i)}$, and $R^{4(i)}$ are independently selected from the group consisting of hydrogen, halogen, hydroxy, trifluoromethyl, cyano, aliphatic hydrocarbyl residue optionally substituted with one or more functional groups and optionally interrupted by one or more heteroatoms, and aromatic hydrocarbyl residue; or $R^{1(i)}$ and $R^{2(i)}$ together form a bridge;

k is 0 or 1;

$A^{(i)}$ and $D^{(i)}$ are independently a saturated or unsaturated optionally substituted aliphatic hydrocarbyl residue, optionally interrupted by a heteroatom or a functional group;

E is a bond or is a heterocyclic residue having one or two ring nitrogen atoms or one ring nitrogen atom and one ring oxygen atom, linked to $D^{(i)}$ and G through a ring nitrogen atom and a ring carbon atom or through two ring nitrogen atoms; and

G is selected from the group consisting of hydrogen, an aliphatic or araliphatic residue, an unsaturated or aromatic monocyclic or polycyclic carbocyclic residue, a saturated, unsaturated, or aromatic monocyclic or polycyclic heterocyclic residue, bonded directly or through a functional group derived from a carbon, nitrogen, oxygen, sulfur, or phosphorus atom,


and the stereoisomers or racemic or non-racemic mixtures of stereoisomers thereof,

C1 *part 1*
 and the tautomers thereof when G is a heterocyclic aromatic ring or an aromatic ring substituted by a hydroxy, mercapto, or amino group,

and the pharmacologically acceptable acid addition salts thereof.

41. (Twice amended) A pharmaceutical composition comprising:

(a) at least one compound selected from the group consisting of compounds of formula I:



where:

each of $R^{1(i)}$, $R^{2(i)}$, $R^{3(i)}$, and $R^{4(i)}$ are independently selected from the group consisting of hydrogen, halogen, hydroxy, trifluoromethyl, cyano, aliphatic hydrocarbyl residue optionally substituted with one or more functional groups and optionally interrupted by one or more heteroatoms, and aromatic hydrocarbyl residue; or $R^{1(i)}$ and $R^{2(i)}$ together form a bridge;

k is 0 or 1;

$A^{(i)}$ and $D^{(i)}$ are independently a saturated or unsaturated optionally substituted aliphatic hydrocarbyl residue, optionally interrupted by a heteroatom or a functional group;

E is a bond or is a heterocyclic residue having one or two ring nitrogen atoms or one ring nitrogen atom and one ring oxygen atom, linked to $D^{(i)}$ and G through a ring nitrogen atom and a ring carbon atom or through two ring nitrogen atoms; and


G is selected from the group consisting of hydrogen, an aliphatic or araliphatic residue, an unsaturated or aromatic monocyclic or polycyclic carbocyclic residue, a saturated, unsaturated, or aromatic monocyclic or polycyclic heterocyclic

residue, bonded directly or through a functional group derived from a carbon, nitrogen, oxygen, sulfur, or phosphorus atom, and the stereoisomers or racemic or non-racemic mixtures of stereoisomers thereof,


and the tautomers thereof when G is a heterocyclic aromatic ring or an aromatic ring substituted by a hydroxy, mercapto, or amino group,

and the pharmacologically acceptable acid addition salts thereof;


(b) at least one compound selected from the group consisting of compounds of formulae II, IIa, IIb, III, IIIa, IIIb, IIIc, IV, IVa, IVb, V, Va, and Vb:




(II)




(IIa)




(IIb)




(III)




(IIIa)




(IIIb)




(IIIc)




(IV)




(IVa)




(IVb)



(V)



(Va)



(Vb)

where:

a is an integer of 1 through 6;

b is an integer of 1 through 2;

X⁻ is selected from the group consisting of fluoride, chloride, bromide, iodide, hydrogensulfate, mesylate, trifluoromethanesulfonate, tosylate, tetrafluoroborate, dihydrogenphosphate, and acetate;

R²¹ is selected from the group consisting of hydrogen, halogen, cyano, alkyl, trifluoromethyl, hydroxyalkyl, hydroxy, alkoxy, alkanoyloxy, alkylthio, aminoalkyl, amino, alkylamino, dialkylamino, formyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, and carboxy;

R²² is selected from the group consisting of hydrogen, halogen, alkyl, trifluoromethyl, hydroxyalkyl, hydroxy, alkoxy, alkanoyloxy, aminoalkyl, amino, alkoxycarbonyl, aminocarbonyl, and carboxy;

12
61
R²³ is selected from the group consisting of hydrogen, alkyl, and hydroxyalkyl;

R²⁴ is selected from the group consisting of alkyl, alkenyl, hydroxyalkyl, alkoxyalkyl, and aralkyl;

R²⁵ is such that the alcohol R²⁵(OH)_a is selected from monovalent linear and branched C₁₋₁₀ alkanols and ω-dialkylaminoalkanols, benzyl alcohol, divalent linear and branched C₂₋₁₀ diols, mono- or divalent C₅₋₇ cycloalkanols, C₅₋₇ cycloalkanediols, C₅₋₇ cycloalkanemethanols, saturated C₅₋₇ heterocyclomethanols, tri-, tetra-, penta-, and hexavalent linear, branched, and cyclic alcohols with 3 to 10 carbon atoms, glycerin, 2,2-bis(hydroxymethyl)-1-octanol, erythritol, pentaerythritol, arabitol, xylitol, sorbitol, mannitol, isosorbitol, tetra(hydroxymethyl)cyclohexanol, and inositol;

12
cont.
R²⁶ is selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, dialkylaminoalkyl, and carboxymethyl;

when b is 1, R²⁷ is selected from the group consisting of hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, dialkylaminoalkyl, and carboxymethyl;

when b is 2, R²⁷ is alkylene in which a methylene group is optionally replaced by O, NH, or N-alkyl; and their thioxo analogs,

and the acid addition salts or anionic salts thereof; and

(c) at least one physiologically acceptable carrier.